Web Images Videos Maps News Shopping Gmail more

Sign in

COOXIC scholar "Rab11A" * "HIV-1"

2005

- 2009

Searc

Scholar All articles Recent articles

Results 1 - 10 of about 45. (0.09 sec)

HIV-1 Vpu promotes release and prevents endocytosis of nascent retrovirus particles from

SJ Neil, SW Eastman, N Jouvenet, PD ... - PLoS Pathog, 2006 - pathogens.plosjournals.org

... A recent study that showed that expression of DN Rab11a or MyosinVb can inhibit

HIV-1 release in the presence of Vpu suggests that a functioning recycling ...

Cited by 63 - Related articles - Cached - All 9 versions

… Recycling Endosome Plays a Key Role in Vpu-mediated Enhancement of **HIV-1** Particle *…*

V Varthakavi, RM Smith, KL Martin, A ... - Traffic, 2006 - pt.wkhealth.com

... In distinct contrast to Vpu, HIV-1 Gag did not appear to accumulate in perinuclear

locations where Vpu and Rab11a overlap but instead was found predominantly ...

Cited by 22 - Related articles - BL Direct - All 4 versions

Rab9 GTPase is required for replication of human immunodeficiency virus type 1, filoviruses

...- ▶ nih.aov

JL Murray s, M Mavrakis, NJ McDonald, M ... - Journal of virology, 2005 - Am Soc Microbiol

... the Rab9 GTPase is required for the unrelated enveloped viruses HIV-1, Marburg virus ... expression of the host genes TIP47, p40, PIKfyve, and Rab11A, indicating a ...

Cited by 48 - Related articles - Bt. Direct - All 11 versions

The cell biology of HIV-1 and other retroviruses

EO Freed, AJ Mouland - Retrovirology, 2006 - biomedcentral.com

... Disruption of recycling endosome function with dominant-negative versions of **Rab11a**

or myosin ... HIV-2 possesses a Vpu-like ability to enhance HIV-1 release; this ...

Cited by 23 - Related articles - Cached - All 10 versions

HIV-1 Vpu inhibits accumulation of the envelope glycoprotein within clathrin-coated, Gag-...

N Van Damme, J Guatelli - Cellular Microbiology, 2008 - interscience.wiley.com

... Vpu itself appears to require transit through rab11a-positive recycling endosomes

to enhance ... vesicles both in macrophages infected with wild-type HIV-1 and in ...

Cited by 5 - Related articles - St. Direct - All 3 versions

Chemokine receptor internalization and intracellular trafficking-

NF Neel, E Schutyser, J Sai, GH Fan, A ... - Cytokine and Growth Factor Reviews, 2005 - Elsevier Cited by 65 - Related articles - All 9 versions

Cellular cofactors involved in HIV assembly and budding

P Spearman - Current Opinion in HIV and AIDS, 2006 - co-hivandaids.com

... that phosphatidylinositol-4,5-bisphosphate [PI(4,5)P 2] regulates HIV-1 Gag targeting

to ... Rab9 and Rab11a were found to be important for HIV replication in one ...

Cited by 1 - Related articles

Host factors exploited by retroviruses

SP Goff - Nature Reviews Microbiology, 2007 - nature.com

... In some cells, notably macrophages, HIV-1 Gag accumulates in intracellular vesicles

126 ... proteins and effectors, notably the proteins Rab9 and Rab11A, in virion ...

Cited by 47 - Related articles - 8L Direct - All 3 versions

... 1 release from latently infected U1 cells reveals a critical role for protein prenylation in HIV-1

T Amet, M Nonaka, MZ Dewan, Y Saitoh, X Qi ... - Microbes and Infection, 2008 - Elsevier

... Rab11a mediates HIV-1 replication in U1 cells 3.5. Rab11a ... View Within Article.

3.4. Rab11a mediates HIV-1 replication in U1 cells. In ...

Cited by 1 - Related articles - All 3 versions

... (shRNA) screening of Jurkat T-cells for human proteins contributory to productive HIV-1 ...-

▶ jbc.org (por)

ML Yeung, L Houzet, VSRK Yedavalli, KT ... - Journal of Biological Chemistry, 2009 - ASBMB

... 1 infection while others are genes whose knock down results in abortive HIV-1 infection. ...

(14), and three other identities (NFKB1, RPL3, and RAB11A) between our ...

Related articles - All 2 versions

Goooogle »

Result Page:

1 2 3 4 5

Next

Rab11A" * "HIV-1"

Search

Go to Google Home - About Google - About Google Scholar

©2009 Google